

# Cross-calibration results from observations of the INS RX J1856, and the WDs HZ 43 and Sirius B



Vadim Burwitz


Intl. Calibration Working Group  
June 14-16 2006, Iceland




# Overview

- Absolute Calibration Soft X-rays
  - dependant on model spectra of WDs
  - what models to use?, uncertainties?
- Analyse LETG + HRC-S data HZ 43A, Sirius B and RX J1856.5-3754
  - determine spectral parameters
  - improve effective area.
- Compare obtained photon spectra with EUVE and ROSAT
  - and XMM (work by Frank Haberl)

# NLTE pure H model spectrum of HZ43




# NLTE pure H model spectrum of HZ43




**Fig. 2.** NLTE pure hydrogen spectra for  $T_{\text{eff}}=50700$  K and  $\log g=8.0$ , calculated with TMAP using two versions of the bound-free and free-free absorption coefficients, Seaton's approximation (dashed curve) and the full Karszas & Latter (1961) description (solid curve). Both models yield practically identical results at optical/ultraviolet wavelengths, but differ by a factor of 1.33 at  $44\text{\AA}$  (vertical dotted line). The insert shows an expanded view. Two further dotted lines mark  $\lambda = 1300\text{\AA}$  and  $\lambda = 5450\text{\AA}$

# HZ43 temperature vs. gravity




**Fig. 3.** Summary of  $T_{\text{eff}}\text{--}\log g$  determinations for HZ43 A from fits to the Balmer lines (circles) and the Lyman lines (triangles and rhombs): Napiwotzki et al. (1993) (Balmer, open circle); Finley et al. (1997) (Balmer, light gray filled circle); Barstow et al. (2003) (Balmer, gray filled circle); this work (STIS Balmer lines, black solid circle); Barstow et al. (2001) (*ORFEUS*, gray lozenge; HUT, open circle); Dupuis et al. (1998) (*ORFEUS*, light gray rhomb); Barstow et al. (2003) (*FUSE* Lyman, gray filled circle); this work (*FUSE* Lyman, black solid triangle); Barstow et al. (1995) (*EUVE* continuum,  $\times$ ); and Vennes & Dupuis (2002) (*EUVE* continuum, asterisk). The error ellipses indicate the results of the present fits to the *Chandra* soft X-ray continuum and refer to the 68%, 90% and 95% confidence levels.

# RX J1856-37 two temperature bb fit




# RX J1856-37 two temperature bb fit




**Fig. 4.** *Chandra* LETG+HRC spectrum of RX J1856 binned to  $0.5\text{\AA}$ , derived with the effective area at wavelengths longwards of the carbon edge corrected to fit an abundance of carbon in the interstellar medium of  $a(C) = 245 - 400$  ppm with 60% in dust grains. The two-temperature fit shown is for 320 ppm, with the upper level of the step indicated for 245 and 400 ppm, too. The fit has a  $\chi^2 = 97.7$  for 115 dof (see Sect. 4.5). The model parameters are quoted in Table 2.

# Simultaneous fit to RXJ1856 and the WDs





# Simultaneous fit to RXJ1856 and the WDs



**Fig. 5.** Simultaneous fit of RXJ1856, HZ43 A, and Sirius B in the wavelength ranges marked by vertical dotted lines (see Sect. 4.4.2). The LETG spectra binned to  $0.5\text{\AA}$  are shown as data points, the corresponding best-fit models as solid curves, and the first-order contributions as dashed curves. The area correction function  $\alpha$  is shown at the top. It converts the nominal LETG+HRC-S first-order effective area  $A^0$  of the November 2004 release into the adjusted area  $A$  used in this paper. Systematic uncertainties in  $\alpha$  are indicated by error bars at 46, 70, 90, and  $125\text{\AA}$ . The steps in the count rate spectra of HZ43 A and RXJ1856 at 49 and  $69\text{\AA}$  result from the detector gaps. Sirius B was observed off axis and its gaps are located differently (see text).

# Parameters obtained from fit


| Parameter  | Value±Error         |
|--|---------------------|
| (a) <b>HZ43 A</b> ( $\lambda = 45 - 160 \text{ \AA}$ )   |                     |
| $T_{\text{eff}}$ (K)                                     | $51126 \pm 660$     |
| $\log g$   | $7.90 \pm 0.08$     |
| $R^2/d^2$ ( $10^{-23}$ )                                 | $3.011 \pm 0.010$   |
| $N_{\text{HI}}$ ( $10^{17} \text{ cm}^{-2}$ )            | $8.91 \pm 0.37$     |
| (b) <b>Sirius B</b> ( $\lambda = 74 - 160 \text{ \AA}$ ) |                     |
| $T_{\text{eff}}$ (K)                                     | $24923 \pm 115$     |
| $\log g$   | $8.6 f^1$           |
| $R^2/d^2$ ( $10^{-21}$ )                                 | $4.877 \pm 0.010$   |
| $N_{\text{HI}}$ ( $10^{17} \text{ cm}^{-2}$ )            | $6.5 \pm 2.0^2$     |
| (c) <b>RX J1856</b> ( $\lambda = 15 - 74 \text{ \AA}$ )  |                     |
| $kT_{\text{spot}}$ (eV)                                  | $62.83 \pm 0.41$    |
| $kT_{\text{star}}$ (eV)                                  | $32.26 \pm 0.72$    |
| $R_1/d$ (km/pc)  | $0.0378 \pm 0.0003$ |
| $R_2/d$ (km/pc)  | $0.1371 \pm 0.0010$ |
| $N_{\text{HI}}$ ( $10^{20} \text{ cm}^{-2}$ )            | $1.10 \pm 0.03$     |

**Table 2.** Parameters of HZ43 A, Sirius B, and RX J1856 based on the simultaneous fit of our model spectra to the LETG+HRC count rate spectra in the wavelength intervals given. The quoted  $1-\sigma$  ( $\Delta\chi^2 = +1$ ) errors are correlated and derived from fits with the other parameters for each object kept free. The letter *f* indicates: fixed.



<sup>1</sup> Based on Barstow et al. (2005); Holberg et al. (1998)

<sup>2</sup> Hébrard et al. (1999). Our fit is required to stay within the  $1-\sigma$  error.


# Comparison of photon spectra



# RX J1856.5-3754: A 'stable' neutron star



Individual fits with  
all parameters free:



Simultaneous fits with  
 $N_H$  linked:

**Model:**  
**tbabs\*body**

from Frank Haberl

# Summary

**Context.** The absolute calibration of space-borne instruments in the soft X-ray regime rests strongly on model spectra of hot white dwarfs.

**Aims.** We analyze the *Chandra* LETG+HRC observations of the white dwarfs HZ43 A and Sirius B and of the neutron star RX J185635–3754 with the aim of resolving current uncertainties in the soft X-ray spectral fluxes and photospheric parameters of the three stars. We apply the derived photon spectra to a cross-calibration of the LETG+HRC-S with the short-wavelength *EUVE* spectrometer and the *ROSAT* PSPC.

**Methods.** We tie HZ43 A to the flux of RXJ1856 in the 44–48Å range and perform a simultaneous least squares fit to the LETG+HRC spectra of the three stars. This allows us to determine an internally consistent set of spectral energy distributions and an empirically derived wavelength-dependent correction to the LETG+HRC-S effective area. We employ NLTE model atmospheres calculated with *TMAP* for the white dwarfs and a two-blackbody model for RXJ1856, tied to the respective optical fluxes.

**Results.** The two-blackbody model for RXJ1856 features a hot spot on a cooler star and yields  $kT_{\text{spot}} = 62.8 \pm 0.4$  eV and  $kT_{\text{star}} = 32.3 \pm 0.7$  eV with a stellar radius as seen from infinity of  $16.0 \pm 0.1$  km for a distance of 117 pc. For HZ43 A, our fit yields  $T_{\text{eff}} = 51126 \pm 660$  K and  $\log g = 7.90 \pm 0.080$  (cgs) with anti-correlated errors ( $1-\sigma$ ) which include not only the statistical but also the systematic uncertainties of the fit. HZ43AB displays a previously detected bremsstrahlung component with a temperature  $kT \simeq 0.6$  keV. For Sirius B, we find  $T_{\text{eff}} = 24923 \pm 115$  K for fixed  $\log g = 8.6$ . The calibration of the short-wavelength *EUVE* spectrometer differs from that of the LETG+HRC-S by  $15 \pm 7\%$ . The *ROSAT* PSPC is found to be correctly calibrated within a few percent and reports of a major miscalibration are unfounded.

**Conclusions.** We have obtained improved parameters for RXJ185635–3754, HZ43 A, and Sirius B which fit the observations from the optical to the soft X-ray regime. Our approach allows us to quote their absolute spectral fluxes at selected wavelengths which may aid the calibration of other space-borne instruments.